

This is the peer reviewed version of the following article: Greenwood, S., Ruiz-Benito, P., Martínez-Vilalta, J., Lloret, F., Kitzberger, T., Allen, C. D., Fensham, R., Laughlin, D. C., Kattge, J., Bönisch, G., Kraft, N. J. B. and Jump, A. S. (2017), Tree mortality across biomes is promoted by drought intensity, lower wood density and higher specific leaf area. *Ecol Lett*, 20: 539–553, which has been published in final form at <https://doi.org/10.1111/ele.12748>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Tree mortality across biomes is promoted by drought intensity, lower wood density and higher specific leaf area

Sarah Greenwood¹, Paloma Ruiz-Benito^{1,2}, Jordi Martínez-Vilalta^{3,4}, Francisco Lloret^{3,4}, Thomas Kitzberger⁵, Craig D. Allen⁶, Rod Fensham^{7,8}, Daniel C. Laughlin⁹, Jens Kattge^{10,11}, Gerhard Bönisch¹⁰, Nathan J. B. Kraft¹² and Alistair S. Jump^{1,3}

¹Biological and Environmental Sciences, University of Stirling, Scotland, FK9 4LA; ² Forest Ecology and Restoration Group, Department of Life Sciences, Science Building, University of Alcalá, Campus Universitario, 28871 Alcalá de Henares, Madrid, Spain; ³CREAF Cerdanyola del Vallès 08193, Spain, ⁴Universidad Autònoma Barcelona, Cerdanyola del Vallès 08193, Spain; ⁵Laboratorio Ecotono, INIBIOMA, CONICET-Universidad Nacional del Comahue, Bariloche, Río Negro, Argentina; ⁶U.S. Geological Survey, Fort Collins Science Center, New Mexico Landscapes Field Station, Los Alamos, New Mexico 87544, USA; ⁷Queensland Herbarium, Environmental Protection Agency, Mt Coot-tha Road, Toowong, Queensland 4066, Australia; ⁸School of Biological Sciences, University of Queensland, St Lucia, Queensland, 4072, Australia; ⁹Environmental Research Institute and School of Science, University of Waikato, Hamilton, New Zealand; ¹⁰Max Planck Institute for Biogeochemistry, Hans-Knöll-Straße 10, 07745 Jena, Germany; ¹¹German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig Deutscher Platz 5e, 04103 Leipzig, Germany; ¹²Department of Ecology and Evolutionary Biology, University of California, Los Angeles, 621 Charles E. Young Drive South, Los Angeles, CA, 90095, USA.

AUTHORSHIP: SG, PRB, JMV, FL, TK, CDA, RF and ASJ conceived and designed the study. SG collected data and performed statistical analyses. DCL, JK, GB and NK contributed data. SG, PRB, JMV, FL, TK, CDA, RF and ASJ wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

Accepted for publication in *Ecology Letters* published by Wiley-Blackwell

DATA ACCESSIBILITY: Should the manuscript be accepted, all relevant data would be archived in the Dryad public repository.

Corresponding author: Sarah Greenwood, Biological and Environmental Sciences,
University of Stirling, Scotland, FK9 4LA, sarah.greenwood@stir.ac.uk,
sarah.greenwood85@gmail.com, +441786467767

Running title: Tree mortality across biomes

Keywords: Climate change, die-off, forest dynamics, functional traits.

Type of article: Reviews and Syntheses

Number of: Words in abstract: 200, Main text: 6714, References: 158, Figures: 6, Tables: 2

Abstract

Drought events are increasing globally, and reports of consequent forest mortality are widespread. However, due to a lack of a quantitative global synthesis, it is still not clear whether drought-induced mortality rates differ among global biomes and whether functional traits influence the risk of drought-induced mortality. To address these uncertainties, we performed a global meta-analysis of 58 studies of drought-induced forest mortality. Mortality rates were modelled as a function of drought, temperature, biomes, phylogenetic and functional groups, and functional traits. We identified a consistent global-scale response, where mortality increased with drought severity (log mortality ($\text{trees trees}^{-1} \text{ year}^{-1}$) increased 0.46 (95% CI=0.2-0.7) with one SPEI unit drought intensity). We found no significant differences in the magnitude of the response depending on forest biomes or between angiosperms and gymnosperms or evergreen and deciduous tree species. Functional traits explained some of the variation in drought responses between species (i.e. increased from 30

to 37% when wood density and specific leaf area were included). Tree species with denser wood and lower specific leaf area showed lower mortality responses. Our results illustrate the value of functional traits for understanding patterns of drought-induced tree mortality and suggest that mortality could become increasingly widespread in the future.

INTRODUCTION

Increased frequency and intensity of drought events, defined as deviation from local long-term average climate, are predicted to occur throughout the world as a consequence of global climate change (Dai, 2013; Trenberth et al. 2014). Forest mortality related to extreme drought has already been reported worldwide (Allen et al. 2010, 2015), with potentially far-reaching impacts on forest diversity and function (Slik, 2004), human society and economies (Anderegg et al. 2013; Hanewinkel et al. 2013), biogeochemical cycling (Bonan 2008; Adams et al. 2010), and species distribution (Benito-Garzon et al. 2013). Forests play a major role in driving global climate due to their key role in hydrological cycling (Bonan 2008; Frank et al. 2015) and ability to sequester and store large amounts of carbon (Pan et al. 2011). Evidence already exists for a widespread reduction in carbon sequestration during drought in tropical (Gatti et al. 2014; Doughty et al. 2015) and temperate forest ecosystems (Anderegg et al. 2015).

There are many examples of drought-induced tree mortality, and several reviews and region-specific syntheses (e.g. Fensham et al. 2009; Allen et al. 2010; Phillips et al. 2010). Additionally, the information gleaned from observational studies of drought-induced tree mortality is supplemented by broad-scale experimental work, which usually allows clearer attribution of causes and deeper mechanistic understanding (Estiarte et al. 2016; Meir et al.

2015). There is however, a lack of quantitative synthesis at a global scale. Despite recent syntheses addressing size-dependent mortality (Bennett et al. 2015) and the importance of hydraulic traits in global-scale mortality (Anderegg et al. 2016), it is difficult to discern broad-scale patterns or to predict future responses. While differential mortality is often reported between co-existing species (e.g. Mueller et al. 2005; Ruthrof et al. 2015), and studies have increased our understanding of the physiological mechanisms of tree mortality (Mencuccini et al. 2015; Davi and Cailleret, 2017), there is still a lack of knowledge of drought-induced mortality responses of different tree functional strategies at a global scale.

Examples of recent mortality related to drought and heat stress exist across all forested continents and major forest types (Figure 1). In Northern temperate forests, high mortality rates are reported for *Pinus sylvestris*, associated with droughts in the 1990s and 2000s (Dobbertin et al. 2005; Galiano et al. 2010) and for *Populus tremuloides* in western Canada (Hogg et al. 2008) and in the USA (Worrall et al. 2008; Ganey and Vojta, 2011; Kane et al. 2014). Drought impacts are particularly well known in the Southwest USA (Allen et al. 2015), with high mortality of numerous tree species, including: *Juniperis monosperma* (Gitlin et al. 2006), *Pinus ponderosa*, *Pinus edulis*, *Pseudotsuga menziesii*, and *Abies concolor* (Ganey and Vojta 2011; Kane et al. 2014). Mortality is also well documented in Mediterranean areas, for example in Spain, high mortality is occurring in *Pinus halepensis* plantations (García de la Serrana et al. 2015), in natural forests in the Sierra Nevada (Herrero et al. 2013) and in Scots pine populations in the Pyrenees and the north east (Vilà-Cabrera et al. 2013). In the southern hemisphere, broad-scale tree die-off events are affecting multiple *Eucalyptus* and *Corymbia* species in Australia (Fensham and Holman, 1999; Rice et al. 2004; Fensham et al. 2009; Matusick et al. 2013) and *Nothofagus* and *Austrocedrus* stands in Patagonia (Suarez and Kitzberger 2008). In tropical areas, extreme droughts associated with

El Niño-Southern Oscillation events have led to increased mortality in species-rich tropical forests in Asia (Aiba and Kitayama, 2002), Central America (Chazdon et al. 2005) and South America (Williamson et al. 2000) while 2005 and 2010 droughts in Amazonia were also associated with mortality increases (Phillips et al. 2009; Feldpausch et al. 2016).

Some studies suggest that drought and/or increased temperature impacts on demography will be greatest in, or limited to, already arid ecosystems (Ruiz-Benito et al. 2014). Indeed, a synthesis by Anderegg et al. (2015) found evidence of stronger reductions in tree growth after drought in dry ecosystems. Conversely, we could expect that the trees in these drier ecosystems may be more adapted to water stress (Arndt et al. 2015), and that the species growing in generally moister conditions could be most sensitive to extreme drought events (Jump et al. in press). Similarly, individuals growing nearer the dry range edge of their species may be more sensitive, but not necessarily, as local adaptation could play a role in counteracting the effects of drier conditions (Lázaro-Nogal et al. 2016). Current uncertainty about which areas are likely to be the most vulnerable to drought is an obstacle for conservation and management planning, while gaps in our understanding of the physiological explanations for drought-induced mortality, and high variation between locations and species make long-term and broad-scale predictions of forest vulnerability and community composition difficult (McDowell et al. 2013; Meir et al. 2015).

Drought-induced tree mortality can lead to long-term changes in the composition and structure of forests due to species and age-specific sensitivities (Martínez-Vilalta and Lloret, 2016). Cailleret et al. (2013) predict future broad-scale mortality of the dominant tree species *Picea abies* and *Abies alba* and their replacement with *Fagus sylvatica* in response to summer droughts in Bavaria, and Ruthrof et al. (2015) show that *Eucalyptus marginata* is more

sensitive to drought than *Corymbia calophylla* in southwestern Australia and predict future compositional shifts. Several examples exist where high mortality of dominant tree species suggests a shift in community composition; montane forests of Switzerland and northern Spain may be undergoing a vegetation shift due to increased drought. *Pinus sylvestris* suffers high mortality and poor regeneration and tends to be replaced by the more drought tolerant *Quercus* species (Galiano et al. 2010; Rigling et al. 2013; Aguade et al. 2015). In the Southwest USA *Pinus edulis* suffers high mortality during drought and *Juniperus* spp are becoming dominant (Mueller et al. 2005), or forest die-off leads to a persistent shift of vegetation type from forest to woodland (Allen and Breshears, 1998). In Patagonia, severe drought events have led to a shift in forest composition, with *Nothofagus dombeyi* being replaced with *Austrocedrus chilensis* (Suarez and Kitzberger, 2008). In extreme cases, changes in drought conditions could potentially lead to a biome shift, resulting in, for example, a seasonally dry tropical forest becoming a savanna, especially when other drivers of change, such as deforestation, act in synergy with drought (Staal et al. 2015).

Changes in tree species and functional composition directly affect forest structure and functioning, although today there is still debate on which functional groups are the most sensitive to drought. Functional groups are composed of individuals with shared physiological responses to, and impacts on, the environment (Tilman et al. 1997). Individuals are often classified into groups based on shared values of particular traits, with the aim of aiding the prediction of response to changes in environment (Hooper and Vitousek, 1997). Evidence exists for differential response of species depending on functional traits and groups; e.g. Phillips et al. (2010) found a higher sensitivity to drought in larger trees with lower-density wood in tropical forests. The functional composition of these forests is, therefore, expected to shift in favour of smaller trees with denser wood, which could result in a

reduction in carbon storage (Stephenson et al. 2014; Fauset et al. 2015). The study of changes in functional composition, and the differential response of species depending on functional traits or broad functional groups, could offer some scope for generalisation of drought response and the mechanisms involved. Indeed, differences in response have been identified based on broad taxonomic and functional groupings; for example, gymnosperms are generally shown to be more sensitive to drought than angiosperms (Anderegg et al. 2015). In Mediterranean and temperate regions of Europe, angiosperms and gymnosperms show contrasting responses to global change. While drying climate reduces growth in both, an increase in temperature and changes in recent management have been associated with increased dominance of angiosperms at the expense of gymnosperms (Gómez-Aparicio et al. 2011; Coll et al. 2013; Rigling et al. 2013; Vayreda et al. 2016). Widespread evidence also exists that larger trees generally may be more at risk from drought-induced mortality (Phillips et al. 2010; Bennett et al. 2015; McDowell and Allen, 2015). The importance of tree size has been suggested in both observational studies and broad-scale experiments in tropical forests (Nepstad et al. 2007; Rowland et al. 2015). Other traits implicated in drought responses, such as wood density (Pratt et al. 2007), specific leaf area (Valladares and Sánchez-Gómez 2006) and root depth (Irvine et al. 2002, 2004) also are likely to affect tree sensitivity to drought.

Given the likely impacts of climate change on ecosystem composition and function, there is an urgent need to synthesise data on tree mortality in order to better understand current patterns and predict the impacts on different forest biomes at a global scale. Here we synthesised data from 58 papers including 398 tree species, from 33 locations and 42 drought events worldwide (Figure 1). We modelled mortality data (proportion of standing dead trees and annual rates) in response to drought and temperature, and we explored differences between biomes and functional groups and investigated the influence of specific functional

traits. Specifically, we ask: 1) Is there a consistent broad-scale relationship between drought, temperature and forest mortality, or are there significant differences between biomes? 2) Are there differences in mortality based on major phylogenetic or functional groups such as angiosperms/gymnosperms or deciduous/evergreen species? 3) Do functional traits explain variation in species-level mortality?

METHODS

A global dataset of forest mortality in response to drought

We sought relevant peer-reviewed publications via Web of Science, Google Scholar and Scopus using combinations of the following search terms: “mortality”, “drought”, “forest”, “tree”, “dieback”, “die-off”, “defoliation”, “drying” and “climate change”. Additional references from the review of Allen et al. (2010) were also included. We screened papers for relevance based on the abstract and saved them for further consideration if they met the general criteria of reporting drought-induced tree mortality. Papers were selected for final inclusion in the meta-analysis if they met the following criteria: 1) mortality data were provided, as a proportion of trees dead, either as a single value based on one survey or as an annualized rate calculated between two or more survey periods; these two types of data were analysed separately. 2) Surveys were ground-based, from aerial photographs or satellite imagery, or a combination of these methods. 3) Mortality was attributed to drought (either one event or a series of events). Experimental studies were only included when the control plots were subject to natural drought during the experimental period and then only the data from these controls were included. 4) The data were collected at the plot level, with a minimum plot size of 0.01 ha, and a minimum number of trees for each species and plot of

five individuals. 5) The survey year(s) and the drought year(s) were clearly reported. 6) Cases where mortality occurred due to fire acting in combination with drought were excluded, but studies where biotic interactions such as attack by bark beetles had occurred were included. Finally, 7) mortality was reported for adult trees (not seedlings or saplings) at the species level. Initially, we studied more than 200 drought and mortality relevant papers in detail, and extracted relevant data. Finally, 58 papers met all the inclusion criteria, 30 of these provide annualized mortality rates, and 28 have single survey proportional values of standing mortality. The case studies cover all forested continents and major forest types (Figure 1). Thirty-three different locations (at the geographic scale of country or state in the case of USA) and 42 drought events (location/time combinations) are covered by our dataset. Selection criteria and particularly the need for quantitative mortality information led to several important and broad-scale studies being excluded (e.g. Phillips et al. 2009).

From each publication, we extracted information on the standing proportional mortality (% dead trees of all trees of each species) or an annual rate of mortality ($m = 1 - (N_1/N_0)^{1/t}$, where N_0 and N_1 are the number of trees present at the beginning and end of the survey interval, t (Sheil et al. 1995)), at the species-level. We report this species-specific mortality as a proportion: trees trees⁻¹ year⁻¹. We also recorded species identity, geographic coordinates, year of the drought, plot size (ha) and sample size (No. plots and trees). If the information was available, we recorded site data as stand density (No. trees ha⁻¹), soil descriptions (e.g. order/classification, depth, texture) and management history. While such site-specific differences will introduce additional variation into the response of trees to drought, these factors were rarely reported, and we could not incorporate them into the analysis. We were also unable to account for forest age or successional stage, as this was

rarely reported. Full information on all studies and species included in the analysis is available in Supporting Information (Appendix S1 and S2).

Drought, temperature and functional trait data at a global scale

In order to assess the relationship between climate change and mortality we collected drought and temperature data. We calculated drought with the Standardised Precipitation Evapotranspiration Index (SPEI), using the mean SPEI at a 12-month timescale, obtained from SPEIbase v.2.3 (2014) (<http://hdl.handle.net/10261/104742>). SPEI is a multi-scalar drought index available as a global gridded dataset at a spatial resolution of 0.5° and has advantages over other drought indices as it allows for the identification of drought at different temporal scales, whilst also being sensitive to changes in evaporative demand (Vicente-Serrano et al. 2010). It has weaknesses, in that soil properties are not accounted for so it does not include a measure of water extractability, but we chose it here based on the evidence that it outperforms other traditional indices such as PDSI (Palmer drought severity index) and SPI (standardised precipitation index) (Vicente-Serrano et al. 2010; Bachmair et al. 2015; Blauhut et al. 2016). SPEI is calculated based on the difference (D) between monthly precipitation and potential evapotranspiration ($D=P-PET$), with the D values aggregated at different timescales from 3 to 24 months, following a procedure whereby SPEI is expressed as a standardised index, with negative values indicating drought over the timescale considered, relative to median values for a long-term reference period. SPEI currently covers the period January 1901 to December 2014. We chose a 12-month period, as it should identify prolonged droughts, of significance for trees. For temperature, we used monthly data from CRUTEM4 dataset (HadCRUT4, <https://crudata.uea.ac.uk/cru/data/temperature/>), a global gridded dataset at 0.5° spatial resolution (Osborn and Jones, 2014) developed by the Climatic

258 Research Unit (University of East Anglia) and the Hadley Centre (UK Met Office). For each
259 study the climate data described above were extracted for the reported geographical
260 coordinates for seven temporal windows: each of 1-5 years preceding the final date of
261 mortality data collection reported in the study, the period between surveys (only for annual
262 rate data), and the reported drought periods. Mean and minimum values of SPEI and mean
263 and maximum values for temperature were calculated from monthly data for each of these
264 periods for every study. We considered the SPEI for up to 5 years before the mortality
265 occurred because a lag is often reported between drought and subsequent mortality. Although
266 these climatic datasets are excellent resources for our global analysis, the somewhat coarse
267 spatial resolution at which they are available potentially reduces the variation that we can
268 explain since we are unable to consistently describe climate at a finer site-specific scale.

269

270 Each study was categorised into a broad biome type based on geographic location, species
271 composition, climate conditions and forest type. Biome types are broadly based on the
272 categories described by Whittaker (1975). We categorised each study to one of the following
273 types: temperate gymnosperm (mid-high latitude and/or altitude forests composed mainly of
274 gymnosperm species); temperate angiosperm (mid-high latitude and/or altitude forests
275 composed mainly of angiosperm species); Mediterranean and dry woodlands (Mediterranean
276 systems and dry temperate, subtropical woodlands and woodland savanna ecotones); tropical
277 (wet tropical forests). While these biome categories encompass diverse forest types,
278 alternative groupings did not lead to different results when the effect of biome was modelled.
279 We obtained information on phylogenetic groups depending on angiosperm or gymnosperm
280 classification and functional groups depending on leaf phenology/lifespan
281 (evergreen/deciduous) by searching the literature for species information in peer reviewed
282 publications. We obtained functional trait data from the TRY traits database (wood density

(WD), specific leaf area (leaf area per unit dry mass; SLA), maximum tree height (MH) (Kattge et al. 2011; Appendix S2)) and from the Royal Botanic Gardens Kew Seed Information Database (seed mass; SM: Royal Botanic Gardens Kew, 2016)). Full trait data was available for 171 of the 398 species in our dataset (see Appendix S2 for species list and data). The subset of species for which trait data were available was similar in terms of the balance between biomes to the full annual mortality dataset (Appendix S3, Table S3.1). For a small number of species, the trait value was extrapolated from the genus level, by averaging the value for three or more species of the same genus (details in Appendix S2). We selected the traits WD, SLA, MH and SM to achieve a balance between drought relevance and data availability. These traits are widely recognised as key traits of plant function and performance; they are relevant to growth, survival and reproduction and can determine species positions along major axes of ecological strategies of resource acquisition and use (Swenson et al. 2012; Díaz et al. 2016). Traits such as P50 (the xylem pressure inducing 50% loss of conductivity; Choat et al. 2012) may be more directly related to drought but hydraulic traits were not available for most species. We show the relationship between WD and SLA and raw mortality (Fig. S3.1 and S3.2 in Appendix S3) and P50 (Figure S3.3). For the species we modelled, we found a weak positive correlation between P50 and WD ($r=0.2$), and a stronger positive correlation between P50 and SLA ($r=0.75$ for a logarithmic fit; Figure S3.3). While the TRY traits database represents an excellent resource, we acknowledge that there are limitations associated with the application of species mean values. Variation in study methods, and differences in sample sizes between species occur for traits deposited in TRY, which could lead to some uncertainty around trait values, and there is no consideration of intraspecific variation. Nevertheless, we do not expect such variability to introduce a systematic bias into our analyses.

Statistical analysis

We followed a linear, three-step modelling process whereby we determined the effects on species-specific tree mortality of: 1) different timescales of SPEI and temperature; 2) biome and phylogenetic and functional groups; and 3) functional traits. We fitted linear mixed effect models using a normal error distribution and log link following an Information Theoretic Approach (Burnham and Anderson, 2002), where models are ranked based on AIC (Akaike Information Criterion) values. Models were fitted with the lme4 package (Bates et al. 2015) in R 3.2.2 (R Core Team 2015). Annualised and proportional species-level mortality data were log transformed prior to analysis, and were analysed separately due to the different nature of these data types. Since there were some zero values in the dataset, 0.001 was added to every mortality value before transformation.

To be sure of the representativeness of the timescale selected, we determined the effects of different timescales of SPEI and temperature on species-level mortality rate. Mortality was modelled against mean and minimum SPEI and mean and maximum temperature calculated at the seven timescales defined (each of 1-5 years before final mortality data collection, year(s) of reported drought, the years between surveys). The predictor variables were standardised to z scores prior to modelling using the scale function in R, which subtracts the mean and divides by the standard deviation. Models containing different combinations of the timescales of SPEI and temperature were compared with the best model selected through ranking of AICc values (AIC corrected for small sample size, Burnham and Anderson 2004; Bunnefeld and Phillimore 2012), and the importance of fixed effects was assessed through the magnitude of parameter estimates. We consider a reduction of AICc of 2 or more to indicate a significantly better model (Burnham and Anderson, 2004).

333 Additionally, a likelihood ratio test was performed to test the significance of the best model
334 (as indicated by AICc value) compared to a null model containing only random effects, and
335 the next best model in the AICc ranking. SPEI and temperature variables were included as
336 fixed effects in all models, and a study-specific identifier was included as a random intercept
337 to account for multiple species occurring in one study and for the site-specific information
338 such as tree density and soil information that we were unable to include in the models. The
339 different timescales of drought were investigated first through model comparison
340 parameterised only with SPEI, and then temperature was added as an additional fixed effect
341 to the model including the optimum drought timescale.

342
343 Once the most parsimonious combination of climatic fixed effects was determined,
344 the differences between biomes and due to major phylogenetic and functional groups were
345 investigated (i.e. angiosperms/gymnosperms and deciduous/evergreen, respectively). Biome
346 was included as a fixed main effect and angiosperm/gymnosperm, deciduous/evergreen
347 groups were included as a fixed main effect and an interaction term with climate (SPEI and
348 temperature). Each model was compared to the same model without the inclusion of the
349 respective subgrouping, based on model AICc values and likelihood ratio test.

350
351 Finally, the effects of functional traits on drought-induced mortality rates were
352 determined. The four continuous functional traits (specific leaf area, maximum height, wood
353 density and seed mass) were added to the best model identified through the process described
354 above as both fixed effects and fixed effect interactions with climate. All traits were initially
355 included and were sequentially removed based on parameter estimates, model AICc values
356 and likelihood ratio tests, as described above. Prior to analysis, we checked for correlation
357 and collinearity between traits. No pair of traits was significantly correlated (maximum r of

0.15 for WD and SLA) or collinear (maximum variance inflation factor (VIF; Dormann et al. 2012) of 1.2). We also used PCA to investigate trait collinearity and to determine if any combination of traits represented a certain functional strategy. No consistent strategies were identified (Table S3.2, Appendix 3). Trait data were standardised to z scores before modelling to allow parameter estimates to be compared. These functional trait models were fitted based on a subset of the data for which the functional trait data were available (171 of 398 species), so AICc values are not comparable with the full optimum model.

To better explore the direction of the important trait interaction effects identified through the above modelling process, predicted values based on models including each trait individually were plotted as 3D surface plots. These 3D plots were created in SigmaPlot (SigmaPlot v. 12.5 Systat software, San Jose); all other plots were created in R 3.2.2. (R Core team 2015).

RESULTS

The available standing proportional mortality data are based on single surveys and are limited in their value because it is not possible to determine the date of death, since the trees standing dead at one point in time may have died over a wide range of dates. Therefore, only the results from the analysis of annualised data, which could be attributed to specific drought events, are presented here. However, the results of this analysis were largely consistent with those for standing proportional mortality. The results for standing proportional mortality data, which were analysed separately, are available in Appendix S4.

Annualised mortality data consisted of 30 studies, 308 species and 27 drought events (Figure 1; Appendix S1). Drought intensity of the reported drought event was highest in Arizona in 2000-2004 with a SPEI value of -3.3 (a monthly mean value; Negron et al. 2009) and lowest in Costa Rica at -0.8 in 1997-1998 (Chazdon et al. 2005). Annual mortality rates (species-level) varied between 0.4 trees trees⁻¹ yr⁻¹ (Negron et al. 2009) and zero (various sources).

Relationship between mortality, drought intensity and temperature

The observed SPEI values for the studies included in our analysis show some divergence from global values. A comparison of the distribution of SPEI associated with the study areas and mortality events, relative to the global values over the full time period covered by the studies (1977-2014), shows more negative SPEI values for the time periods associated with reported mortality events (the 5 years before final mortality surveys). There is, however, a large degree of overlap in the distributions of SPEI for the study period and the five years preceding the reported mortality event (Appendix S5).

The mean SPEI of the reported drought years was the best climatic predictor of annual mortality (Table 1). We identified a consistent response to drought that occurred across all biomes, with mortality increasing as values of SPEI decreased, i.e. as drought, defined as deviation from a long term mean, became more intense (Figure 2a, b). There is little support for the inclusion of biome as a fixed interaction with drought ($\Delta AICc = 3.2$) and the parameter estimates for biome comparisons suggests that the difference in mortality between biomes is small (Figure 3).

Functional groups, phylogenetic groups and functional trait effects on drought-induced tree mortality

We detected no significant difference in the response to drought between angiosperms and gymnosperms or deciduous and evergreen species. The inclusion of these groups as fixed-effect interaction terms with SPEI did not improve the model (AICc increased by 0.2 and 0.7, respectively) and parameter estimates suggest no difference in drought mortality response (Figure 4a, b).

The functional traits WD and SLA influenced the drought-induced mortality response; the inclusion of these traits improved the model compared to one that included drought effects alone (Table 2; Figure 5a, b). However, MH and SM did not correlate with drought responses (Table 2). Drought-induced mortality was higher among species with lower values of WD (Figure 6a) and higher values of SLA (Figure 6b). We observed that the slope of interaction with WD was highest at very negative values of SPEI (more severe drought). For the interaction with SLA it is stronger at higher values of SPEI, (i.e. when drought is less severe), and the relationship is reversed compared to that seen at more negative values of SPEI, where mortality is slightly lower when SLA is higher (Figure 6b).

DISCUSSION

Global-scale patterns of drought-induced mortality

We found a consistent response of tree mortality to drought across biomes, indicative of a global-scale threat to forests from drought, which is consistent with the widespread nature of

recent reports of forest mortality (e.g. Allen et al. 2010, 2015). The fact that we find evidence for a linear increase in log (mortality) (Figures 2 and 5a) with increasing drought intensity means that the relationship with SPEI is exponential and, therefore, increases in drought could have critical consequences for forests. This result is indicative of a threshold response, and contrasts with other broad-scale studies that have suggested a linear relationship between mortality and drought (e.g. Phillips et al 2010). Such a response at the species-level seems realistic, given what we know of responses to drought intensity in individual studies (e.g. Meir et al. 2015). Note, however, that the shape of the drought-mortality response is likely to differ depending on the metric used to characterize drought intensity.

Our study is the first to identify a global-scale effect of drought intensity on tree mortality at the species-level; previous syntheses have either been limited geographically (Fensham et al. 2009; Phillips et al. 2010), or have focused on drivers such as size-dependence of mortality (Bennett et al. 2015) and the influence of hydraulic traits (Anderegg et al. 2016), rather than global patterns. It has been suggested that only dry forests are at risk from drought-induced mortality (Steinkamp and Hickler, 2015). Studies of tree growth have shown responses to drought occurring across the northern hemisphere, although with differences in timescale of response depending on long-term average aridity (Vicente-Serrano et al. 2014). Recent work provides a physiological basis for this global-scale vulnerability to drought-induced mortality, indicating that some 70% of forest species operate at narrow hydraulic safety margins irrespective of biome (Choat et al. 2012). Therefore, species in both arid and mesic habitats could be at high risk of embolism and related mortality if drought increases.

Our synthesis suggests that there is little difference between angiosperms and gymnosperms or between evergreen and deciduous species in the response of mortality to

drought, in line with a recent synthesis by Anderegg et al. (2016). Our global-scale analysis complements that of Anderegg et al. (2016) as these authors use a strict meta-analytic approach by comparing species within a study and consider hydraulic traits, while we focus on a range of traits that are more available and have broad applicability because they relate to multiple aspects of plant functioning (Díaz et al. 2016). We also account explicitly for drought intensity through the inclusion of SPEI data in our models. The fact that we obtain some common results, despite the different methods applied, gives increased support to both analyses. Although our selection of traits allowed for a large sample size, given the greater availability of data, there are certain drawbacks to consider, as the traits we study are unlikely to be directly related to drought-mortality responses. The use of traits with a stronger mechanistic basis (e.g., hydraulic traits) for which the relationship with drought-induced mortality is more direct is probably preferable when they are available.

While the global scale of our study is valuable for identifying broad-scale patterns, we acknowledge that the range of species and biomes included may have led to a masking of differences between groups, since in some cases the advantages of one functional strategy in one biome may not be applicable in another. For example, to be an evergreen species has different implications in temperate *versus* tropical forests. Furthermore, the small sample size and associated statistical limited power also may have constrained our ability to find differences between biomes and functional and phylogenetic groups. Despite these potential limitations, the global patterns that we identify and their congruence with related work by other authors demonstrates the value of this work for increasing our understanding of forest drought risk worldwide.

Functional traits mediate drought response

482

483 Our results suggest that high wood density can be associated with lower susceptibility to
484 drought. Although wood density is a complex trait that may be associated with many
485 ecological signals (Broderson, 2016; Gleason et al. 2016) this finding is substantially in
486 accordance with reported functionality of this trait. Wood properties correlated with density
487 affect water storage and transport, because the width and length of conduits determines
488 hydraulic conductance (Chave et al. 2009) and wood density affects mechanical stability and
489 is associated with resistance to drought-induced cavitation (Hacke et al. 2001; Pratt et al.
490 2007). In a study of saplings and seedlings from dry tropical forest, Markesteijn et al. (2011)
491 found a lower vulnerability to cavitation (an ability to withstand more negative P50) in
492 species with higher wood density. They also found evidence for a trade-off between
493 cavitation resistance and hydraulic conductivity, suggesting that the ability to withstand
494 drought leads to a competitive disadvantage during times when water is not limiting.
495 However, weak support was found for this trade-off in a global synthesis by Gleason et al.
496 (2016), which showed that many species can have both low hydraulic conductivity and low
497 cavitation resistance, and that high wood density is not necessarily associated with low
498 conductivity and high cavitation resistance. Generally, species with high wood density are
499 associated with lower growth rates and lower background mortality rates (Chave et al. 2009).
500 The results of our study show that this strategy can lead to lower mortality rates when
501 exposed to drought stress. In contrast to our findings, Hoffmann et al. (2011) found that
502 although species with higher density wood had higher cavitation resistance, their mortality
503 under drought was higher, due to an inability to prevent declines in water potential, compared
504 to species with less dense wood. Hoffmann et al. studied the response to an especially severe
505 drought at the community level, whereas we consider a wide range of drought conditions and
506 vegetation types, suggesting that differences in overall drought severity and spatial scale

might explain the differences between the results that they report and those presented here. It is notable that despite the complexity of this trait and its relationship with hydraulic performance, we still find a significant, although noisy, global signal.

Our analyses provide evidence that tree species with high specific leaf area can be more susceptible to drought-induced mortality than species with lower specific leaf area, although the nature of the relationship varied with drought severity. Specific leaf area determines plant response to changing resource availability (i.e. water and light), photosynthetic potential and growth rates (Poorter et al. 2009). Depending on stomatal properties, higher specific leaf area could imply a higher potential for leaf water loss; in contrast, a low specific leaf area is often associated with xeromorphic plants, adapted to high water stress (Bussotti et al. 2015). Furthermore, woody species associated with arid niches are characterised by relatively lower specific leaf area in the Mediterranean (Costa-Saura et al. 2016). An experimental study by Valladares and Sánchez-Gómez (2006) showed within-species lower specific leaf area if seedlings were exposed to drought, although across species, a high specific leaf area was associated with greater drought tolerance. Reducing specific leaf area under prolonged drought conditions seems to be consistent across species and studies (Poorter et al. 2009) and is suggestive of a higher resistance to drought in plants with a lower specific leaf area. Although we identify a global-scale signal in the relationship between drought and SLA, as for wood density, many ecological signals are associated with SLA making this relationship highly complex. In the future, the integration of directly relevant traits such as P50 could improve the approach taken here.

Several studies have shown that larger trees could have lower resistance to drought (e.g. Bennett et al. 2015). For example, an experimental drought in a Brazilian Amazonian

rainforest resulted in much larger increases in the mortality rates of larger trees compared to smaller ones (Nepstad et al. 2007; Rowland et al. 2015) and a pan-tropical analysis also revealed that larger trees tend to suffer most during drought (Phillips et al. 2010). A global-scale synthesis by Bennett et al. (2015) shows that drought-related mortality is generally greater in larger trees; they analysed drought mortality related to DBH, but they hypothesised that the relationship is due to height effects. However, in our global-scale analysis that explicitly included species maximum height, we did not find a strong effect due to tree height. This apparent contrast might be because the importance of the other traits that we considered overwhelmed any effect of height, but it also could be due to the different approach taken in our study compared to that of Bennett et al. Our analysis was based on species-level maximum height, while their study used size classes at the plot level irrespective of species. Future work should seek to incorporate individual-level tree heights and stand age, as this will provide the most direct assessment on the importance of tree size.

Implications

Our results suggest that at a global scale, trees with lower wood density and higher specific leaf area tend to be more sensitive to drought. Future increases in the intensity and duration of drought events could therefore lead to changes in forest composition and structure, with implications for forest diversity and ecosystem function. Changes in functional diversity due to the drought sensitivity of different species will also affect forest functioning and drought resilience (Laureto et al. 2015). Forests are an important global carbon sink (Pan et al. 2011) and changes in forest energy exchange could have major consequences for global climate (Frank et al. 2015). Wood density and specific leaf area are both traits that are important for carbon sequestration (Prado-Junior et al. 2016) and storage in aboveground biomass (Chave

et al. 2014). It is, therefore, important to consider the impacts that the greater vulnerability associated with particular values of these traits under drought conditions could have on broad-scale carbon storage (Finegan et al. 2015). Generally, higher biomass might be expected to be stored in ecosystems rich in species with levels of these traits associated with high growth rates (low wood density and high SLA (Grime 1998)), but this is context dependent. For example, in dry tropical forests, species with high wood density and low specific leaf area are associated with high growth rates and carbon storage, precisely because they can continue to function during drought (Prado-Junior et al. 2016). Our results also have successional implications; early successional tree species are characterised by low wood density and high specific leaf area (Aiba et al. 2016). If disturbance increases due to drought-induced mortality, then these early successional species could dominate forests, making the community more susceptible to future die-off.

Our synthesis identifies global-scale drought-induced tree mortality responses, and our analyses of species-specific mortality rates with climate and functional trait data provide quantitative insights into global-scale patterns. However, our best model only explained 37% of the variation in mortality rate. To reliably assess the impact of drought events, and to predict the consequences for forests in the future, effective long-term monitoring across a wide range of forest types is required. In addition, detailed reporting of site conditions would allow future syntheses to consider the importance of factors such as tree density in drought-induced mortality at a global scale (Jump et al. in press). Here, we identified drought-induced mortality responses worldwide that are species-specific, but further analysis depending on stand and tree level characteristics are critical to adequately identify and manage vulnerable forests. Additionally, the consideration of intra-specific variation in functional traits could significantly improve our ability to predict the responses of forests to future drought. The

traits we have identified as critical for drought resistance at a global scale are known to vary within species across environmental gradients (Fajardo and Piper, 2011), thus trait variation across environmental gradients (e.g. Vilà-Cabrera et al. 2015) should be incorporated in future models to better predict climate change impacts in forests worldwide. Although our results do not have high predictive power in themselves, they provide an advance, since our current ability to predict drought-induced mortality is limited (Meir, Mencuccini and Dewar, 2015) and this prediction needs global assessment. Our study highlights the global vulnerability of forests to drought-induced tree mortality and shows that widely available functional traits, such as wood density and specific leaf area, can improve explanatory power.

Conclusion

Based on a synthesis of studies of drought-induced tree mortality events at a global scale we find evidence of greater forest mortality risk from more severe droughts, consistent across biomes and major functional groups. Using functional traits, we could explain an additional fraction of the variation in drought-induced tree mortality. Trees with less dense wood and high specific leaf area may be most at risk. These results further illustrate the value of species-level trait data for understanding general trends in species responses. Overall, our findings indicate that, with increasingly severe future droughts projected, associated drought-induced tree mortality could become increasingly widespread, with significant implications for forest diversity, ecosystem function and climate feedbacks.

ACKNOWLEDGEMENTS

This research was supported by The Leverhulme Trust via International Network grant IN-2013-004, together with the European Union Seventh Framework Programme under PCOFUND-GA-2010-267243 (Plant Fellows) co-funded by the University of Stirling. The study has been supported by the TRY initiative on plant traits (<http://www.try-db.org>). The

TRY initiative and database is hosted, developed and maintained by J. Kattge and G. Bönisch (Max Planck Institute for Biogeochemistry, Jena, Germany). TRY is/has been supported by DIVERSITAS, IGBP, the Global Land Project, the UK Natural Environment Research Council (NERC) through its program QUEST (Quantifying and Understanding the Earth System), the French Foundation for Biodiversity Research (FRB), and GIS “Climat, Environnement et Société” France. Support to CD Allen provided by the U.S. Geological Survey's Ecosystems and Climate & Land Use Change mission areas (through the Western Mountain Initiative). We also thank the following authors for kindly providing us with their original tree mortality data: Shin-Ichiro Aiba, Robin Chazdon, Joseph Ganey, Eric Gustafson and Íñigo Grazow de la Cerda. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

REFERENCES

- Aakala, T., Kuuluvainen, T., Wallenius, T. & Kauhanen, H. (2011). Tree mortality episodes in the intact *Picea abies*-dominated taiga in the Arkhangelsk region of northern European Russia. *J. Veg. Sci.*, 22, 322-33.
- Adams, H. D., Macalady, A. K., Breshears, D. D., Allen, C. D., Stephenson, N. L., Saleska, S. R., Huxman, T. E. & McDowell, N. (2010). Climate-Induced Tree Mortality: Earth System Consequences, *Eos Trans. AGU*, 91, 153–154.
- Aguade, D., Poyatos, R., Rosas, T. & Martínez-Vilalta, J. (2015) Comparative Drought Responses of *Quercus ilex* L. and *Pinus sylvestris* L. in a Montane Forest Undergoing a Vegetation Shift. *Forests*, 6, 2505-2529.
- Aiba, S. I. & Kitayama, K. (2002). Effects of the 1997–98 El Niño drought on rain forests of Mount Kinabalu, Borneo. *J. Trop. Ecol.*, 18, 215-230

639 Aiba, M., Kurokawa, H., Onoda, Y., Oguro, M., Nakashizuka, T. & Masaki, T. (2016).
640 Context-dependent changes in the functional composition of tree communities along
641 successional gradients after land-use change. *J. Ecol.* doi:10.1111/1365-2745.12597
642

643 Allen, C.D. & Breshears, D.D. (1998). Drought-induced shift of a forest-woodland ecotone:
644 rapid landscape response to climate variation. *PNAS*, 95, 14839-14842.
645

646 Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M.,
647 Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E.H., Gonzalez, P., Fenhsam, R., Zhang,
648 Z., Castro, J., Demidova, N., Lim, J-H., Allard, G., Running, S.W., Semerci, & Cobb, N.
649 (2010) A global overview of drought and heat-induced tree mortality reveals emerging
650 climate change risks for forests. *For. Ecol. Man.*, 259, 660-684.
651

652 Allen, C. D., Breshears, D. D & McDowell, N.G. (2015). On underestimation of global
653 vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene.
654 *Ecosphere*, 6, 129.
655

656 Anderegg, W. R. L., Berry, J. A., Smith, D. D., Sperry, J. S., Anderegg, L. D. L. & Field, C.
657 B. (2012) The roles of hydraulic and carbon stress in a widespread climate-induced forest
658 die-off. *PNAS*, 109, 233-237.
659

660 Anderegg, W.R.L., Kane, J.M. & Anderegg, L.D.L. (2013). Consequences of widespread tree
661 mortality triggered by drought and temperature stress. *Nat. Clim. Change*, 3, 30-36.
662

663 Anderegg, W.R.L., Schwalm, C., Biondi, F., Camarero, J.J., Koch, G., Litvak, M., Ogle, K.,
664 Shaw, J.D., Shevliakova, E., Williams, A.P., Wolf, A., Ziaco, E. & Pacala, S. (2015)

665 Pervasive drought legacies in forest ecosystems and their implications for carbon cycle
666 models. *Science*, 349, 528-532.

667

668 Anderegg, W.R.L., Klein, T., Bartlett, M., Sack, L., Pellegrini, A.F.A., Choat, B. & Jansen,
669 S. (2016). Meta-analysis reveals that hydraulic traits explain cross-species patterns of
670 drought-induced tree mortality across the globe. *PNAS*, doi: 10.1073/pnas.1525678113

671

672 Anderson-Teixeira, K.J., Miller, A.D., Mohan, J.E., Hudiburg, T.W., Duval, B.D. & De Lucia,
673 E.H. (2013). Altered dynamics of forest recovery under a changing climate. *Glob. Chan.*
674 *Biol.*, 19, 2001-2021.

675

676 Arndt, S.K., Sanders, G.J., Brsitow, M., Hutley, L.B., Beringer, J. & Livesley, S.J. (2015).
677 Vulnerability of native savanna trees exotic *Khaya senegalensis* to seasonal drought. *Tree*
678 *Phys.*, 35, 783-791.

679

680 Bachmair, S., Kohn, I & Stahl, K. (2015). Exploring the link between drought indicators and
681 impacts. *Nat. Hazards Earth Syst. Sci.*, 15, 1381–1397.

682

683 Bates, D., Maechler, M., Bolker, B., Walker, S. (2015). Fitting Linear Mixed-Effects Models
684 Using lme4. *J. Statistical Software*, 67, 1-48.

685

686 Becker P., Lye, C.O & Goh F. (1998). Selective drought mortality of dipterocarp trees: No
687 correlation with timber group distributions in Borneo. *Biotropica*, 30, 666-671.

688

Benito-Garzón, M., Ruiz-Benito, P. & Zavala, M. A. (2013). Interspecific differences in tree growth and mortality responses to environmental drivers determine potential species distributional limits in Iberian forests. *Glob. Ecol. Biogeog.*, 22, 1141–1151.

Bennett, A.C., McDowell, N.G., Allen, C.D., & Anderson-Teixeira, K.J. (2015). Larger trees suffer most during drought in forests worldwide. *Nature Plants*, 1, 15139, doi: 10.1038/NPLANTS.2015.139

Blauhut, V., Stahl, K., Stagge, J. H., Tallaksen, L. M., De Stefano, L. & Vogt, J. (2016). Estimating drought risk across Europe from reported drought impacts, drought indices, and vulnerability factors. *Hydrol. Earth Syst. Sci.*, 20, 2779–2800,

Bonan, G.B. (2008). Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefit of Forests. *Science*, 320, 1444-1449.

Bowker, M. A., Munoz, A., Martinez, T. & Lau, M. K. (2012). Rare drought-induced mortality of juniper is enhanced by edaphic stressors and influenced by stand density. *J. Arid Environ.*, 76, 9-16.

Brodersen, C.R. 2016. Finding support for theoretical tradeoffs in xylem structure and function. *New Phyt.*, 209, 8–10

Bucci, S.J., Goldstein, G., Meinzer, F.C., Scholz, F.G., Franco, A.C. & Bustamante, M. (2004). Functional convergence in hydraulic architecture and water relations of tropical savanna trees: from leaf to whole plant. *Tree Phys.*, 24, 891-899.

715 Bunnefeld, N. & Phillimore, A.B. (2012). Island, archipelago and taxon effects: mixed
 716 models as a means of dealing with the imperfect design of nature's experiments. *Ecography*
 717 35, 15-22.
 718
 719 Burnham, K. P. & Anderson, D. R. (2002). Model Selection and Multimodel Inference: a
 720 Practical Information-theoretic Approach. Springer.
 721
 722 Burnham, K. P. & Anderson, D. R. (2004). Multimodel Inference: Understanding AIC and
 723 BIC in Model Selection. *Sociological Methods Research*, **33**, 261-304
 724
 725 Bussotti, F., M. Pollastrini, et al. (2015). Functional traits and adaptive capacity of European
 726 forests to climate change. *Environ. Experim. Bot.*, 111, 91-113.
 727
 728 Cailleret, M., Nourtier, M., Amm, A., Durand-Gillmann, M. & Davi, H. (2013). Drought-
 729 induced decline and mortality of silver fir differ among three sites in Southern France. *Ann.*
 730 *For. Sci.*, 71, 643-657.
 731
 732 Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G. & Zanne, A. E. (2009).
 733 Towards a worldwide wood economics spectrum. *Ecol. Lett.*, 12, 351-366.
 734
 735 Chave, J., Rejou-Mechain, M., Burquez, A, Chidumay, E., Colgan, M.S., Delitti, W.B.C. et
 736 al. (2014). Improved allometric models to estimate the aboveground biomass of tropical trees.
 737 *Glob. Chan. Biol.*, 20, 3177-3190.
 738

739 Chazdon, R.L., Brenes, A.R. & Alvarado, B.V. (2005). Effects of climate and stand age on
740 annual tree dynamics in tropical second-growth rain forests. *Ecology*, 86, 1808-1815.
741

742 Choat, B., Jansen, S., Brodribb, T. J., Cochard, H., Delzon, S., Bhaskar, R et al. (2012).
743 Global convergence in the vulnerability of forests to drought. *Nature*, 491, 752-756.
744

745 Coll, M., Peñuelas, J., Ninyerola, M., Pons, X. & Carnicer, J. (2013). Multivariate effect
746 gradients driving forest demographic responses in the Iberian Peninsula. *For. Ecol. Man.*,
747 303, 195-209.
748

749 Condit, R., Hubbell, S.P. & Foster, R.B. (1995). Mortality Rates of 205 Neotropical Tree and
750 Shrub Species and the Impact of a Severe Drought. *Ecol. Mon.*, 65, 419-439
751

752 Costa-Saura, J. M., J. Martínez-Vilalta, Trabucco, A., Spano, D. & Mereu, S. (2016). Specific
753 leaf area and hydraulic traits explain niche segregation along an aridity gradient in
754 Mediterranean woody species. *Per. Plan. Ecol. Evo. Sys.*, 21, 23-30.
755

756 da Costa, A. C. L., Galbraith, D., Almeida, S., Portela, B. T. T., da Costa, M., de Athaydes
757 Silva Junior, J., Braga, A. P. Goncalves, P. H. L., de Oliveira, A. AR., Fisher, R., Phillips, O.
758 L., Metcalfe, D. B., Levy, P., Meir, P. (2010). Effect of 7 yr of experimental drought on
759 vegetation dynamics and biomass storage of an eastern Amazonian rainforest. *New Phyt.*,
760 187,579-591.
761

762 Dai, A. (2013). Increasing drought under global warming in observations and models. *Nat.*
763 *Clim. Change*, 3, 52-58.

764

765 Davi, H & Cailleret, M. (2017). Assessing drought-driven mortality trees with physiological
766 process-based models. *Agr. Forest. Meteorol.*, 232, 279-290.

767

768 Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S. et al. (2016). The
769 global spectrum of plant form and function. *Nature*, 529, 167-171.

770

771 Dobbertin M., Mayer P., Wohlgemuth T., Feldmeyer-Christe E., Graf U., Zimmermann N.E.
772 & Rigling A. (2005). The decline of *Pinus sylvestris* L. forests in the Swiss Rhone valley - A
773 result of drought stress? *Phyton*, 45, 153-156.

774

775 Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carre, G. et al. (2012).

776 Collinearity: a review of methods to deal with it and a simulation study evaluating their
777 performance. *Ecography*, 36, 27-46.

778

779 Doughty, C.E., Metcalfe, D.B., Girardin, C.A.J., Farfan-Amezquita, F., Galiano Cabrera, D.,
780 Huaraca Huasco, W., Silva-Espejo, J.E., Araujo-Murakami, A., da Costa, M.C., Rocha, W.,
781 Feldpausch, T.R., Mendoza, A.L.M., da Costa, A.C.L., Meir, P., Phillips, O.L., & Malhi, Y.
782 (2015). Drought impact on forest carbon dynamics and fluxes in Amazonia. *Nature*, 519, 78-
783 82.

784

785 Estiarte, M., Vicca, S., Peñuelas, J., Bahn, M., Beier, C. Emmett, B.A., Fay, P.A., Hanson,
786 P.J., Hasibeder, R., Kigel, J., Kroel-Dulay, G., Larsen, K.S., Lellei-Kovacs, E., Limousin,
787 J.M., Ogaya, R. Ourcival, J.M., Reinsch, S., Sala, O.E., Schmidt, I.K., Sternberg, M.,
788 Tielborger, K., Tietema, A. & Janssens, I.A. (2016) Few multiyear precipitation-reduction

789 experiments find a shift in the productivity-precipitation relationship. *Glob. Chang. Biol.*, 22,
790 2570-2581.

791

792 Faber-Langendoen, D. & Tester, J.R. (1993). Oak mortality in sand savannas following
793 drought in east-central Minnesota. *Bulletin of the Torrey Botanical Club*, 120, 248-256

794

795 Fajardo, A., Piper, F.I. (2011). Intraspecific trait variation and covariation in a widespread
796 tree species (*Nothofagus pumilio*) in southern Chile. *New Phyt.*, 189, 259-271

797

798 Fan, Z., Fan, X., Crosby, M.K., Moser, W.K., He, H., Spetich, M.A. & Shifley, S.R. (2012).
799 Spatio-temporal trends of oak decline and mortality under periodic regional drought in the
800 Ozark Highlands of Arkansas and Missouri. *Forests*, 3, 614-631.

801

802 Fauset, S., Johnson, M. O., Gloor, M., Baker, T. R., Monteagudo, A., Brien, R. J. W. et al.
803 (2015). Hyperdominance in Amazonian forest carbon cycling. *Nature Comm.*, 6, 6857.

804

805 Feldpausch, T. R., Phillips, O. L., Brien, R. J. W., Gloor, E., Lloyd, J. et al.
806 (2016). Amazon forest response to repeated droughts. *Global Biogeochem. Cycles*, 30, 964–
807 982.

808

809 Fensham, R.J. (1998). The influence of cattle grazing on tree mortality after drought in
810 savanna woodland in north Queensland. *Austral Ecology*, 23, 405-407.

811

812 Fensham, R.J. & Holman, J.E. (1999). Temporal and spatial patterns in drought-related tree
813 dieback in Australian savanna. *J. Appl. Ecol.*, 36, 1035-1050.

814

815 Fensham, R.J., Fairfax, R.J., Ward, D.P. (2009). Drought-induced tree death in savanna.

816 *Glob. Chan. Biol.*, 15, 380-387.

817

818 Finegan, B., Pena-Claros, M., Oliveira, A., Ascarrunz, N., Sydonia Bret-Harte, M., Carreno-

819 Rocabado, G. et al. (2015). Does functional trait diversity predict above-ground biomass and

820 productivity of tropical forests? Testing three alternative hypotheses. *J. Ecol.*, 103, 191-201.

821

822 Floyd, M.L., Clifford, M., Cobb, N.S., Hanna, D., Delph, R., Ford, P. & Turner, D. (2009).

823 Relationship of stand characteristics to drought-induced mortality in three Southwestern

824 piñon - juniper woodlands. *Ecol. App.*, 19, 1223-1230.

825

826 Foden, W., Midgley, G.F., Hughes, G., Bond, W.J., Thuiller, W., Hoffman, M.T., Kaleme, P.,

827 Underhill, L.G., Rebelo, A., Hannah, L. (2007). A changing climate is eroding the

828 geographical range of the Namib Desert tree *Aloe* through population declines and dispersal

829 lags. *Div. Dist.*, 13, 645-653.

830

831 Frank, D.C., Poulter, B., Saurer, M., Esper, J., Huntingford, C., Helle, G., Treydte, K.,

832 Zimmermann, N.E., Schleser, G.H. & Ahlstrom, A. (2015). Water-use efficiency and

833 transpiration across European forests during the Anthropocene. *Nat. Clim. Chan.*, 5, 579-583.

834

835 Galiano, L., Martínez-Vilalta, J & Lloret, F. (2010). Drought-Induced Multifactor Decline of

836 Scots Pine in the Pyrenees and Potential Vegetation Change by the Expansion of Co-

837 occurring Oak Species. *Ecosystems*, 13, 978-991

838

839 Ganey, J. L. & Vojta, S. C. (2011). Tree mortality in drought-stressed mixed-conifer and
840 ponderosa pine forests, Arizona, USA. *For. Ecol. Man.*, 261, 162-168.
841

842 Garcia de la Serrana, R., Vilagrosa, A. & Alloza, J. A. (2015). Pine mortality in southeast
843 Spain after an extreme dry and warm year: interactions among drought stress, carbohydrates
844 and bark beetle attack. *Trees*, 29, 1791-1804.
845

846 Gatti, L.V., Gloor, M., Miller, J.B., Doughty, C.E., Malhi, Y., Domingues, L.G., Basso, L.S.,
847 Martinewski, A., Correia, C.S.C., Borges, V.F., Freitas, S., Braz, R., Anderson, L.O., Rocha,
848 H., Grace, J., Phillips, O.L., Lloyd, J. (2014). Drought sensitivity of Amazonian carbon
849 balance revealed by atmospheric measurements. *Nature*, 506, 76-80.
850

851 Gitlin A.R., Sthultz C.M., Bowker M.A., Stumpf S., Paxton K.L., Kennedy K., Munoz A.,
852 Bailey J.K. & Whitham T.G. (2006). Mortality gradients within and among dominant plant
853 populations as barometers of ecosystem change during extreme drought. *Con. Bio.*, 20, 1477-
854 1486.
855

856 Gleason, S. M., Westoby, M., Jansen, S., Choat, B., Hacke, U. G. *et al.* (2016), Weak tradeoff
857 between xylem safety and xylem-specific hydraulic efficiency across the world's woody plant
858 species. *New Phyt.*, 209, 123–136.
859

860 Gómez-Aparicio, L.,García-Valdés, R., Ruiz-Benito, P. & Zavala, M. A. (2011).
861 Disentangling the relative importance of climate, size and competition on tree growth in
862 Iberian forests: implications for forest management under global change. *Glob. Chan. Biol.*,
863 17, 2400-2414.

864

865 Granzow-de la Cerda, I., Lloret, F., Ruiz, J. E. & Vandermeer, J. H. (2012). Tree mortality
866 following ENSO-associated fires and drought in lowland rain forests of Eastern Nicaragua.
867 *For. Ecol. and Man.*, 265, 248-257.

868

869 Grime, J.P. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder
870 effects. *J. Ecol.*, 86, 902–910.

871

872 Gu, L., Pallardy, S.G., Hosman, K.P. & Sun, Y. (2015) Predictors and mechanisms of the
873 drought-influenced mortality of tree species along the isohydric to anisohydric continuum in a
874 decade-long study of a central US temperate forest. *Biogeosciences Discussions*, 12, 1285-
875 1325.

876

877 Hacke, U.G., Sperry, J.S., Pockman, W.T., Davis, S.D. & McCulloh, K.A (2001). Trends in
878 wood density and structure are linked to prevention of xylem implosion by negative pressure.
879 *Oecologia*, 126, 457-461.

880

881 Hanewinkel, M., Cullmann, D.A., Schelhaas, M-J., Nabuurs, G-J., Zimmermann, N.E. (2013)
882 Climate change may cause severe loss in the economic value of European forest land. *Nat.*
883 *Clim. Chan.*, 3, 203-307.

884

885 Heitzman, E., Grell, A., Spetich, M. & Starkey, D. (2007). Changes in Forest Structure
886 Associated with Oak Decline in Severely Impacted Areas of Northern Arkansas. *South. J.*
887 *Appl. For*, 31, 17-22.

888

889 Herrero, A., Castro, J., Zamora, R., Delgado-Huertas, A. & Querejeta, J. I. (2013). Growth
 890 and stable isotope signals associated with drought-related mortality in saplings of two
 891 coexisting pine species. *Oecologia*, 173, 1613-1624.
 892
 893 Hoffmann, W.A., Marchin, R.M., Abit, P. & Lau, O.L. (2011). Hydraulic failure and tree
 894 dieback are associated with high wood density in a temperate forest under extreme drought.
 895 *Glob. Chan. Biol.*, 17, 2731-2742.
 896
 897 Hogg, E.H., Brandt, J.P. & Michaelin, M. (2008). Impacts of a regional drought on the
 898 productivity, dieback and biomass of western Canadian aspen forests. *Can. J. For. Res.*, 38,
 899 1373-1384.
 900
 901 Hooper, D.U. & Vitousek, P.M. (1997). The effects of plant composition and diversity on
 902 ecosystem processes. *Science*, 277, 1302–1305.
 903
 904 Huang, C., Asner, G. P., Barger, N. N., Neff, J. C. & Floyd, M. L. (2010). Regional
 905 aboveground live carbon losses due to drought-induced tree dieback in piñon–juniper
 906 ecosystems. *Remote Sensing of Environment*, 114, 1471-1479.
 907
 908 Huang, Q., Swatantran, A., Dubayah, R. & Goetz, S.J. (2014). The influence of vegetation
 909 height heterogeneity on forest and woodland bird species richness across the United States.
 910 *PLoS ONE*, 9, e103236.
 911
 912 Irvine J., Law B.E., Anthoni P.M. & Meinzer F.C. (2002). Water limitations to carbon
 913 exchange in old-growth and young ponderosa pine stands. *Tree Phys.*, 22, 189–196.

914

915 Irvine J., Law B.E., Kurpius M.R., Anthoni P.M., Moore D. & Schwarz P.A. (2004). Age-
916 related changes in ecosystem structure and function and effects on water and carbon
917 exchange in ponderosa pine. *Tree Phys.*, 24, 753–763.

918

919 Itoh, A., Nanami, S, Harata, T., Ohkubo, T., Tan, T., Chong, L., Davies, S.J. & Yamakura, T.
920 (2012). The effect of habitat association and edaphic conditions on tree mortality during El
921 Niño-induced drought in a Bornean dipterocarp forest. *Biotropica*, 44, 606-617.

922

923 Jump, A. S., Ruiz-Benito, P., Greenwood, S., Allen, C. D., Kitzberger, T., Fensham, F.,
924 Martínez-Vilalta, J. & Lloret, F. (2017). Structural overshoot of tree growth with climate
925 variability and the global spectrum of drought-induced forest dieback. *Global Change*
926 *Biology* (in press).

927

928 Kane, J. M., Kolb, T. E., McMillin, J. D. (2014). Stand-scale tree mortality factors differ by
929 site and species following drought in southwestern mixed conifer forests. *For. Ecol. Man.*,
930 330, 171-182.

931

932 Kardol, P., Todd, D.E., Hanson, P.J., & Mulholland, P.J. (2010). Long-term successional
933 forest dynamics: species and community responses to climatic variability. *J. Veg. Sci.*, 21,
934 627-642.

935

936 Kattge, J., Díaz, S., Lavorel, S., et al. (2011) TRY – a global database of plant traits. *Glob.*
937 *Chan. Biol.*, 17, 2905–2935.

938

939 Khan J.A., Rodgers W.A., Johnsingh A.J.T. & Mathur P.K. (1994). Tree and shrub mortality
 940 and debarking by sambar *Cervus unicolor* (Kerr) in Gir after a drought in Gujarat, India. *Biol.*
 941 *Cons.*, 68, 149-154.

942

943 Klos, R.J., Wang, G.G., Bauerle, W.L. & Rieck, J.R. (2009). Drought impact on forest
 944 growth and mortality in the southeast USA: an analysis using forest health and monitoring
 945 data. *Ecol. App.*, 19, 699-708.

946

947 Koepke, D. F., Kolb, T. E. & Adams, H. D. (2010). Variation in woody plant mortality and
 948 dieback from severe drought among soils, plant groups and species within a northern Arizona
 949 ecotone. *Oecologia*, 163, 1079-1090.

950

951 Kukowski, K. R., Schwinning, S. & Schwartz, B.F. (2013). Hydraulic responses to extreme
 952 drought conditions in three co-dominant tree species in shallow soil over bedrock. *Oecologia*,
 953 171, 819-830.

954

955 Kuusinen, M. & Siitonen, J. (1998) Epiphytic lichen diversity in old-growth and managed
 956 *Picea abies* stands in southern Finland. *J. Veg. Sci.*, **9**, 283-292.

957

958 Laureto, L. M. O., Cianciaruso, M. V. & Menezes Samia, D. S. (2015). Functional diversity:
 959 an overview of its history and applicability. *Nat. Conserv.*, 13, 112-116.

960

961 Lázaro-Nogal, A., Matesanz, S., Hallik, L., Krasnova, A., Traveset, A. & Valladares, F.
 962 (2016). Population differentiation in a Mediterranean relict shrub: the potential role of local
 963 adaptation for coping with climate change. *Oecologia*, 180, 1075-1090.

964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987

Linares, J. C., Taiqui, L., Camarero, J. J. (2011). Increasing drought sensitivity and decline of Atlas cedar (*Cedrus atlantica*) in the Moroccan Middle Atlas Forests. *Forests*, 2, 777-796.

Lwanga, J.S. (2003). Localized tree mortality following the drought of 1999 at Ngogo, Kibale National Park, Uganda. *African J. Ecol.*, 41, 194-196.

MacGregor S.D. & O'Connor T.G. (2002). Patch dieback of *Colophospermum mopane* in a dysfunctional semi-arid African savanna. *Austral Ecology*, 27, 385-395.

Markalas, S. (1992). Site and stand factors related to mortality rate in a fir forest after a combined incidence of drought and insect attack. *For. Ecol. Man.*, 47, 367-374.

Markestijn, L., L. Poorter, Paz, H., Sack, L. & Bongers, F. (2011). Ecological differentiation in xylem cavitation resistance is associated with stem and leaf structural traits. *Plant, Cell Envir.*, 34, 137-148.

Martin, P.A., Newton, A.C., Cantarello, E. & Evans, P. (2015). Stand dieback and collapse in a temperate forest and its impact on forest structure and biodiversity. *For. Ecol. Man.*, 358, 130-158.

Martínez-Vilalta, J. & Pinol, J. (2002). Drought-induced mortality and hydraulic architecture in pine populations of the NE Iberian Peninsula. *For. Ecol. Man.*, 161, 247-256.

988 Martínez-Vilalta, J. and Lloret, F. (2016). Drought-induced vegetation shifts in terrestrial
989 ecosystems: the key role of regeneration dynamics. *Glob. Planet. Change*, 144, 94-108.
990

991 Matusick, G., Ruthrof, K. X., Brouwers, N. C., Dell, B. & Hardy, G. St. J. (2013). Sudden
992 forest canopy collapse corresponding with extreme drought and heat in a Mediterranean-type
993 eucalypt forest in southwestern Australia. *European Journal of Forest Research*, 132, 497-
994 510.
995

996 McDowell, N. G., Fisher, R. A., Xu, C., Domec, J. C., Hölttä, T. et al. (2013). Evaluating
997 theories of drought-induced vegetation mortality using a multimodel experimental
998 framework. *New Phyt.*, 200, 304-321.
999

1000 McDowell, N. & Allen, C.D. (2015). Darcy's law predicts widespread forest mortality under
1001 climate warming. *Nature Climate Change*, 5, 669-672.
1002

1003 Meir, P., Mencuccini, M. & Dewar, R. C. (2015). Drought-related tree mortality: addressing
1004 the gaps in understanding and prediction. *New Phyt*, 207, 28-33.
1005

1006 Meir, P., Wood, T.E., Galbraith, D.R., Brando, P.M., Da Costa, A.C.L., Rowland, L. &
1007 Ferreira, L.V. (2016). Threshold responses to soil moisture deficit by trees and soil in tropical
1008 rain forests: insights from field experiments. *Bioscience* 65, 882-892.
1009

1010 Mencuccini, M., Minunno, F., Salmon, Y., Martínez-Vilalta, J. & Hölttä, T. (2015).
1011 Coordination of physiological traits involved in drought-induced mortality of woody plants.
1012 *New Phyt*, 208, 396-409.

1013

1014 Millar, C. I., Westfall, R. D., Delany, D. L., Bokach, M. J., Flint, A. L. & Flint, L. E. (2012).

1015 Forest mortality in high-elevation whitebark pine (*Pinus albicaulis*) forests of eastern

1016 California, USA; influence of environmental context, bark beetles, climatic water deficit, and

1017 warming. *Can. J. For. Res.*, 42, 749-765.

1018

1019 Mokria, M., Gebrekirstos, A., Aynekulu, E. & Brauning, A. (2015). Tree dieback affects

1020 climate change mitigation potential of a dry Afromontane forest in northern Ethiopia. *For.*

1021 *Ecol. Man.*, 344, 73-83.

1022

1023 Mueller, R.C., Scudder, C.M., Porter, M.E., Talbot Trotter III, R., Gehring, C.A. & Whitham,

1024 T.G. (2005). Differential tree mortality in response to severe drought: Evidence for long-term

1025 vegetation shifts. *J. Ecol.*, 93, 1085-1093.

1026

1027 Nakagawa, M., Tanaka, K., Nakashizuka, T., Ohkubo, T., Kato, T., Maeda, T., Sato, K.,

1028 Miguchi, H., Nagamasu, H., Ogino, K., Teo, S., Hamid, A.A. & Seng, L.H. (2000). Impact of

1029 severe drought associated with the 1997–1998 El Niño in a tropical forest in Sarawak. *J.*

1030 *Trop. Ecol.*, 16, 355-367

1031

1032 Negron, J.F., McMillin, J.D., Anhold, J.A., & Coulson, D. (2009). Bark beetle-caused

1033 mortality in a drought-affected ponderosa pine landscape in Arizona, USA. *For. Ecol. Man.*,

1034 257, 1353-1362.

1035

1036 Nepstad, D.C., Tohver, I.M., David, R., Moutinho, P. & Cardinot, G. (2007). Mortality of
 1037 large trees and lianas following experimental drought in an amazon forest. *Ecology*, 88, 2259-
 1038 2269.

1039

1040 Newbery, D.M. & Lingenfelder, M. (2004). Resistance of a lowland rain forest to increasing
 1041 drought intensity in Sabah, Borneo. *J. Trop. Ecol.*, 20, 613-624.

1042

1043 Newbery, D.M. & Lingenfelder, M. (2008). Plurality of tree species responses to drought
 1044 perturbation in Bornean tropical rain forest. *Plant Ecol.*, 201, 147-167.

1045

1046 Olano, J.M. & Palmer, M.W. (2003). Stand dynamics of an Appalachian old-growth forest
 1047 during a severe drought episode. *For. Ecol. Man.*, 174, 139-148.

1048

1049 Osborn, T.J. & Jones, P.D. (2014). The CRUTEM4 land-surface air temperature data set:
 1050 construction, previous versions and dissemination via Google Earth. *Earth System Science*
 1051 *Data*, 6, 61-68.

1052

1053 Pan, Y., Birdsey, R.A., Fang, J., Houghton, R., Kauppi, P.E., Kurz, W.A., Phillips, O.L.,
 1054 Shvidenko, A., Lewis, S.L., Canadell, J.G., Ciais, P., Jackson, R.B., Pacala, S., McGuire,
 1055 A.D., Piao, S., Rautiainen, A., Sitch, S. & Hayes, D. (2011). A large and persistent carbon
 1056 sink in the world's forests. *Science*, 333, 988-993.

1057

1058 Pangle, R. E., Limousin, J-M., Plaut, J. A., Yopez, E. A., Hudson, P. J., Boutz, A. L., Gehres,
 1059 N., Pockman, W. T. & McDowell, N. G. (2015). Prolonged experimental drought reduces

1060 plant hydraulic conductance and transpiration and increases mortality in a pinon-juniper
 1061 woodland. *Ecology and Evolution*, **5**, 1618-1638.

1062

1063 Phillips, O. L., Aragão, L. E. O. C., Lewis, S. L., Fisher, J. B., Lloyd, J. & López-González,
 1064 G. (2009). Drought sensitivity of the Amazon rainforest. *Science*, 323, 1344-1347.

1065

1066 Phillips, O.L., van der Heijden, G., Lewis, S.L., Lopez-Gonzalez, G., Aragão, L.E.O.C.,
 1067 Lloyd, J. et al. (2010). Drought-mortality relationships for tropical forests. *New Phyt.*, 187,
 1068 631-646.

1069

1070 Poorter, H., Niinemets, Ü., Poorter, L. Wright, I.J. & Villar, F. (2009). Causes and
 1071 consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phyt.*, 182,
 1072 565-588.

1073

1074 Potts, M.D. (2003). Drought in a Bornean everwet rain forest. *J. Ecol.*, 91, 467-474.

1075

1076 Poulos, H.M. (2014). Tree mortality from a short-duration freezing event and a global-
 1077 change-type drought in a Southwestern pinon-juniper woodland, USA. *PeerJ*, 2, e404.

1078

1079 Prado-Junior, J.A., Schiavini, I., Vale, V.S., Arantes, C.S., van der Sande, M.T., Lobeck, M.,
 1080 & Poorter, L. (2016). Conservative species drive biomass productivity in tropical dry forests.
 1081 *J. Ecol.*, 104, 817-827.

1082

1083 Pratt, R.B., Jacobsen, A.L., Ewers, F.W. & Davis, S.D. (2007). Relationships among xylem
 1084 transport, biomechanics and storage in stems and roots of nine Rhamnaceae species of the
 1085 California chaparral. *New Phyto.*, 174, 787–798.

1086

1087 R Core Team. (2015). R: A language and environment for statistical computing. R
 1088 Foundation for Statistical Computing, Vienna, Austria.

1089

1090 Rice, K.J., Matzner, S.L., Byer, W. & Brown, J.R. (2004). Patterns of tree dieback in
 1091 Queensland, Australia: the importance of drought stress and the role of resistance to
 1092 cavitation. *Oecologia*, 139, 190–198.

1093

1094 Rigling, A., Bigler, C., Eilmann, B., Feldmeyer-Christe, E., Gimmi, U., Ginzler, C., Graf, U.,
 1095 Mayer, P., Vacchiano, G., Weber, P., Wohlgemuth, T., Zweifel, R. & Dobbertin, M. (2013).
 1096 Driving factors of a vegetation shift from Scots pine to pubescent oak in dry Alpine forests.
 1097 *Glob. Chan. Biol.*, 19, 229–240.

1098

1099 Rowland, L., da Costa, A.C.L., Galbraith, D.R., Oliveira, R.S., Binks, O.J., Oliveira, A.A.R.,
 1100 Pullen, A.M., Doughty, C.E., Metcalfe, D.B., Vasconcelos, S.S., Ferreira, L.V., Malhi, Y.,
 1101 Grace, J., Mencuccini, M. & Meir, P. (2015). Death from drought in tropical forests is
 1102 triggered by hydraulics not carbon starvation. *Nature*, 528, 119–121.

1103

1104 Royal Botanic Gardens Kew. (2016) Seed Information Database (SID). Version 7.1.
 1105 Available from: <http://data.kew.org/sid/> (April 2016).

1106

1107 Ruiz-Benito, P., Madrigal-Gonzalez, J., Ratcliffe S., Coomes, D.A., Kandler, G., Lehtonen,
 1108 A., Wirth, C. & Zavala, M.A. (2014). Stand structure and recent climate change constrain
 1109 stand basal area change in European forests: a comparison across boreal, temperate and
 1110 Mediterranean biomes. *Ecosystems*, 17, 1439-1454.
 1111
 1112 Ruthrof, K. X., Matusick, G., Hardy, G. E. St. J. (2015). Early differential responses of co-
 1113 dominant canopy species to sudden and severe drought in a Mediterranean-climate type
 1114 forest. *Forests*, 6, 2082-2091.
 1115
 1116 Santiago, L.S., Goldstein, G., Meinzer, F.C., Fisher, J.B., Machado, K., Woodreiff, D. &
 1117 Jones, T. (2004). Leaf photosynthetic traits scale with hydraulic conductivity and wood
 1118 density in Panamanian forest canopy trees. *Oecologia*, 140, 543-550.
 1119
 1120 Savage, M. (1997) The role of anthropogenic influences in a mixed-conifer forest mortality
 1121 episode. *J. Veg. Sci.*, 8, 95-104.
 1122
 1123 Sheil, D., Burslem, D.F.R.P. & Alder, D. (1995). The interpretation and misinterpretation of
 1124 mortality rate measures. *J. Ecol.*, 83, 331-333.
 1125
 1126 Slik, J.W. (2004). El Niño droughts and their effects on tree species composition and
 1127 diversity in tropical rain forests. *Oecologia*, 141, 114-120.
 1128
 1129 Smith, J. M., Paritsis, J., Veblen, T. T. & Chapman, T. B. (2015). Permanent forest plots
 1130 show accelerating tree mortality in subalpine forests of the Colorado Front Range from 1982
 1131 to 2013. *For. Ecol. Man.*, 341, 8-17.

1132

1133 Staal, A., Dekker, S.C., Hirota, M. & van Nes, E.H. (2015). Synergistic effects
 1134 of drought and deforestation on the resilience of the south-eastern Amazon rainforest.
 1135 *Ecological complexity* 22, 65-67.

1136

1137 Steinkamp, J. & Hickler, T. (2015) Is drought-induced forest dieback globally increasing? *J.*
 1138 *Ecol*, **103**, 31-43.

1139

1140 Stephenson, N.L., Das, A.J., Condit, R., Russo, S.E., Baker, P.J., Beckman, N.G., Coomes,
 1141 D.A., Lines, E.R., Morris, W.K., Ruger, N. et al. (2014). Rate of tree carbon accumulation
 1142 increases continuously with tree size. *Nature*, 507, 90-93.

1143

1144 Suarez, M.L. & Kitzberger, T. (2008). Recruitment patterns following a severe drought: long-
 1145 term compositional shifts in Patagonian forests. *Can. J. For. Res.*, 38, 3002-3010.

1146

1147 Swenson, N. G., Enquist, B. J., Pither, J. Kerkhoff, A J., Boyle, B. *et al.* (2012) The
 1148 biogeography and filtering of woody plant functional diversity in North and South America.
 1149 *Glob. Ecol. Biogeogr.*, 21, 798-808.

1150

1151 Tafangenyasha, C. (1997). Tree loss in the Gonarezhou National Park (Zimbabwe) between
 1152 1970 and 1983. *Journal of Environmental Management*, 49, 355-366.

1153

1154 Tilman, D., Knops, J., Wedin, D., Reich, P., Ritchie, M. & Siemann, E. (1997). The influence
 1155 of functional diversity and composition on ecosystem processes. *Science*, 277, 1300–1302.

1156

1157 Trenberth, K. E., A. Dai, G. van der Schrier, P. D. Jones, J. Barichivich, K. R. Briffa, & J.
1158 Sheffield, 2014: Global warming and changes in drought. *Nat. Clim. Chan.*, 4, 17-22.
1159

1160 Tsopelas, P., Angelopoulos, A., Economou, A., Voulala, M., & Xanthopoulou, E.,
1161 2001. Monitoring crown defoliation and tree mortality in the fir-forest of Mount Parnis,
1162 Greece. In: Radoglou, K. (ed.) Proceedings of the International Conference: Forest Research,
1163 a Challenge for an Integrated European Approach. August 27-1 September 2001,
1164 Thessaloniki, Greece. pp. 253-258
1165

1166 Tsopelas, P., Angelopoulos, A., Economou, A. & Soulioti, N. (2004) Misteltoe (*Viscum*
1167 *album*) in the fir forest of Mount Parnis, Greece. *For. Ecol. Man.*, 202, 59-65.
1168

1169 Twidwell, D., Wonkka, C. L., Taylor, C. A., Zou, C. B., Twidwell, J. J. & Rogers, W. E.
1170 (2014). Drought-induced woody plant mortality in an encroached semi-arid savanna depends
1171 on topographic factors and land management. *Applied Vegetation Science*, **17**, 42-52.
1172

1173 Valladares, F. & Sánchez-Gómez, D. (2006). Ecophysiological traits associated with drought
1174 in Mediterranean tree seedlings: individual responses versus interspecific trends in eleven
1175 species. *Plant Biol.*, 8, 688-697.
1176

1177 van Nieuwstadt, M.G.L., & Shiel, D. (2005). Drought, fire and tree survival in a Borneo rain
1178 forest, East Kalimantan, Indonesia. *J. Ecol.*, 93, 191-201.
1179

1180 Vayreda, J., Martínez-Vilalta, J., Gracia, M., Canadell, J. G. & Retana, J. (2016).
 1181 Anthropogenic-driven rapid shifts in tree distribution lead to increased dominance of
 1182 broadleaf species. *Glob Change Biol*, 22, 3984–3995.
 1183
 1184 Vicente-Serrano, S.M., Begueria, S., & Lopez-Moreno, J.I. (2010). A Multiscalar Drought
 1185 Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration
 1186 Index. *J. Clim.*, 23, 1696-1718.
 1187
 1188 Vilà-Cabrera, A., Martínez-Vilalta, J., Galiano, L. & Retana, J. (2013). Patterns of forest
 1189 decline and regeneration across Scots pine populations, *Ecosystems*, 16, 323-335.
 1190
 1191 Vilà-Cabrera, A., Martínez-Vilalta, J. & Retana, J. (2015). Functional trait variation
 1192 across environmental gradients in Temperate and Mediterranean trees. *Glob. Ecol.*
 1193 *Biogeog.*, 24, 1377-1389.
 1194
 1195 Viljoen, A.J. (1995). The influence of the 1991/92 drought on the woody vegetation of
 1196 the Kruger National Park. *Koedoe*, 38, 85-97.
 1197
 1198 Vincke, C., Diedhiou, I., Grouzis, M. (2010). Long-term dynamics and structure of
 1199 woody vegetation in the Ferlo (Senegal). *J. Arid Env.*, 74, 268-276.
 1200
 1201 Whittaker, R. H. (1975). *Communities and Ecosystems*. Macmillan, New York.
 1202

1203 Williamson, G. B., Laurance, W. F., Oliveira, A. A., Delamonica, P., Gascon, C., Lovejoy, T.
 1204 E & Pohl, L. (2000). Amazonian tree mortality during the 1997 El Niño drought. *Cons. Bio.*,
 1205 14, 1538-1542.
 1206
 1207 Worrall, J., Egeland, L., Eager, T., Mask, R.A., Johnson, E.W., Kemp, P. & Shepperd, W.D.
 1208 (2008). Rapid mortality of *Populus tremuloides* in southwestern Colorado, USA. *For. Ecol.*
 1209 *Man.*, 255, 686-696.
 1210
 1211 Wright, S.J., Kiajima, K., Kraft, N.J.B., Reich, P.B., Wright, I.J., Bunker, D.E., Condit, R.,
 1212 Dalling, J.W., Davies, S.J., Diaz, S., Engelbrecht, B.M.J., Harms, K.E., Hubbell, S.P., Marks,
 1213 C.O., Ruiz-Jaen, M.C., Salvador, C.M. & Zanne, A.E. (2007). Functional traits and the
 1214 growth-mortality trade-off in tropical trees. *Ecology*, 91, 3664-3674.
 1215
 1216 Zegler, T. J., Moore, M. M., Fairweather, M. L., Ireland, K. B. & Fule, P. Z. (2012). *Populus*
 1217 *tremuloides* mortality near the southwestern edge of its range. *For. Ecol. Man.*, 282, 196-207.
 1218
 1219
 1220 **SUPPORTING INFORMATION**
 1221 **Appendix S1:** Information on the studies and species included in the meta-analysis
 1222 **Appendix S2:** Species list, mean trait values, A/G and D/E classification and sources of trait
 1223 data, including citations and details of the species for which data was extrapolated from the
 1224 genus level.
 1225 **Appendix S3:** Trait data information: biome representation, raw mortality against traits,
 1226 relationship between wood density and specific leaf area and P50, TRY trait reference list.
 1227 **Appendix S4:** Model results from single survey proportional mortality data analysis.

Appendix S5: SPEI distribution of study areas compared to global values over the same period.

TABLES

Table 1: Model comparison using AICc values (AIC corrected for small sample size) and parameter estimates for sets of models used to identify the optimum model(s). The 95% confidence intervals associated with the parameter estimates are shown for the fixed effects in the best model, highlighted in bold. SPEIm represents mean SPEI and TEMPm mean temperature. Confidence intervals were calculated with boot strapped resampling. The random effect included in these models was a study specific identifier. Results of likelihood ratio tests (P-values) are shown for the best models compared to the null model (one including only random effects) and the next best model, as indicated by AICc values.

Fixed effect(s)	Parameter estimate(s) (95% CI)	AICc	Δ AICc	R ² marginal/ conditional	P-value
SPEIm drought years	-0.46 (-0.7 to - 0.2)	1876.7	0.0	0.06/0.29	0.0005; <0.0001
SPEIm between survey years	-0.43	1880.1	3.4	0.05/0.32	
SPEIm drought + TEMPm 3 years pre survey end	-0.46; -0.01	1880.3	3.6	0.06/0.30	
SPEIm drought + TEMPm 5 years pre survey end	-0.46; 0.002	1880.3	3.6	0.06/0.30	

SPEIm drought + TEMPm 2 years pre survey end	-0.46; -0.007	1880.3	3.6	0.06/0.30
SPEIm drought + TEMPm 4 years pre survey end	-0.46; 0.0009	1880.3	3.6	0.06/0.30
SPEIm drought + TEMPm between survey years	-0.46; 0.01	1880.4	3.7	0.06/0.30
Min SPEI drought years	-0.45	1880.5	3.8	0.06/0.34
Min SPEI 2 years pre survey end	-0.54	1880.8	4.1	0.03/0.33
SPEIm + maximum temperature 2 years pre survey end	-0.46; -0.04	1880.8	4.1	0.06/0.30
SPEIm + maximum temperature 3 years pre survey end	-0.46; -0.01	1880.9	4.2	0.06/0.30
SPEIm + maximum temperature 4 years pre survey end	-0.45; 0.01	1880.9	4.2	0.06/0.30
SPEIm + maximum temperature 5 years pre survey end	-0.45; 0.01	1880.9	4.2	0.06/0.30
SPEIm drought + maximum temperature between survey years	-0.45; 0.009	1880.9	4.2	0.06/0.30
Minimum SPEI between	-0.39	1881.5	4.8	0.04/0.34

survey years				
Min SPEI 3 years pre survey	-0.32	1883.0	6.3	0.03/0.33
end				
Min SPEI 4 years pre survey	-0.30	1883.3	6.6	0.03/0.32
end				
Min SPEI 5 years pre survey	-0.31	1883.3	6.6	0.03/0.33
end				
SPEIm 2 years pre survey end	-0.24	1883.8	7.1	0.02/0.31
SPEIm 3 years pre survey end	-0.27	1884.3	7.6	0.02/0.30
SPEIm 5 years pre survey end	-0.19	1885.8	9.1	0.01/0.31
SPEIm 4 years pre survey end	-0.15	1887.0	10.3	0.006/0.31

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257 Table 2: Model comparison and selection when considering functional traits and drought
 1258 response. AICc values (AIC corrected for small sample size) and parameter estimates for sets
 1259 of models used to identify the optimum model. SPEIm represents mean SPEI. The model
 1260 containing the best combination of fixed effects is shown in bold. The traits considered are
 1261 wood density (WD), specific leaf area (SLA), maximum height (MH) and seed mass (SM).
 1262 The results of likelihood ratio tests are shown for the best model (P-value), compared to a
 1263 null and the next best model. Here the null model for comparison contains random effects and
 1264 the fixed effect of mean SPEI of the drought.
 1265

Fixed effect(s)	Parameter estimate(s)	AICc	Δ AICc	R ² marginal/conditional	P-value
SPEIm	-1.28 (-2.04 to	1013.5	0.0	0.14/0.37	0.001;
drought +	-0.52)				0.01
SPEIm	0.38 (-0.2 to				
drought x WD	0.97)				
+ SPEIm	-0.41 (-0.95 to				
Drought x SLA	0.13)				
SPEIm drought	-1.19; 0.39	1014.8	1.3	0.10/0.33	
+ SPEIm					
drought x WD					
SPEIm drought	-1.22; -0.38	1015.7	2.2	0.09/0.34	
+ SPEIm					
drought x SLA					
SPEIm drought	-1.14	1016.6	3.1	0.06/0.30	

SPEIm drought	-1.26; 0.26; -	1016.6	3.1	
+ SPEIm	0.63			
drought x WD				
+ SPEIm				
drought x MH				
SPEIm drought	-1.28; 0.27; -	1016.8	3.3	0.15/0.37
+ SPEIm	0.26; -0.53			
drought x WD				
+ SPEIm				
drought x SLA				
+ SPEIm				
drought x MH				
SPEIm drought	-1.24; -0.22; -	1019.7	6.2	0.10/0.33
+ SPEIm	0.56			
drought x SLA				
+ SPEIm				
drought x MH				
SPEIm drought	-1.28; 0.37; -	1022.8	9.3	0.16/0.38
+ SPEIm	0.27; -0.5; -			
drought x (WD,	0.21			
SLA, MH, SM)				
SPEIm drought	-1.17; -0.12;	1022.9	9.4	0.07/0.31
+ SPEIm	0.02			
drought x SM				

FIGURES

Figure 1: Locations of the studies included in the analysis. Dark grey shading indicates studies providing annual rates of mortality based on two or more surveys and light grey standing proportional mortality values based on a single survey (these are analysed separately with annual mortality rate analyses presented in the main text and single survey proportions presented in supporting information). The biome category of each study is indicated by shape. Temperate gymnosperms: diamonds, temperate angiosperms: triangles, Mediterranean and dry woodlands: circles and tropical: squares. Annual mortality studies: Aiba and Kitayama, 2002; Chazdon et al. 2005; Condit et al 1995; da Costa et al. 2010; Dobbervin et al. 2005; Faber-Langendone and Tester 1993; Fan et al 2012; Ganey and Vojta 2011; Granzow de la Cerda et al. 2012; Gu et al 2015; Hogg et al. 2008; Huang et al. 2010; Itoh et al. 2012; Kane et al. 2014; Kardol et al. 2010; Klos et al. 2009; Kukowski et al. 2013; Mueller et al. 2005; Nakagawa et al. 2000; Negron et al. 2009; Newbery and Lingenfelder 2008; Olano and Palmer, 2003; Potts, 2003; Poulos, 2014; Smith et al. 2015; Tafangenyasha, 1997; Tsopelas, 2001; Tsopelas et al. 2004; van Nieuwstadt and Shiel, 2005; Worall et al. 2008. Single survey proportional mortality studies: Aakala et al. 2011; Becker et al. 1998; Bowker et al. 2012; Cailleret et al. 2013; Fensham, 1998; Fensham and Holman, 1999; Floyd et al. 2009; Foden et al. 2007; Garcia de la Serrana et al. 2015; Gitlin et al. 2006; Heitzman et al. 2007; Khan et al. 1994; Koepke et al. 2010; Linares et al. 2011; Lwanga, 2003; Markalas, 1992; Martínez Vilalta and Pinol, 2002; Matusick et al. 2013; Millar et al. 2012; Mokria et al. 2015; Rice et al. 2004; Savage, 1997; Slik, 2004; Suarez and Kitzberger, 2008; Twidwell et al. 2014; Viljoen, 1995; Vincke et al. 2010; Zegler et al. 2012.

Figure 2: Annual observed mortality data at species-level plotted against mean SPEI for the reported drought years (a), and log-scale annual observed mortality against mean SPEI for the reported drought year, including the model prediction using the parameter estimate for the best model with 95% confidence intervals (b). More negative values of SPEI indicate more severe drought.

Figure 3: Parameter estimates for the model including mean SPEI of the reported drought years and biome as fixed effects to test if drought induced mortality varies by biome. The parameter estimates show the difference in the mortality response to drought of each biome with 95% confidence intervals, calculated using bootstrapped resampling. Sample sizes (number of studies) by biome are as follows: temperate gymnosperms: five, temperate angiosperms: eight, Mediterranean and dry woodlands: seven and tropical: ten.

1303 Figure 4: Parameter estimates for the models including mean SPEI of the reported drought
1304 years and the groups angiosperms/gymnosperms (a) and evergreen/ deciduous (b) as fixed
1305 effect interactions with mean SPEI. 95% confidence intervals are shown, calculated using
1306 bootstrapped resampling.

1307

1308 Figure 5: Relationship between predicted annual mortality and drought (solid line) with 95 %
1309 confidence (dotted lines) based on the best model including the effect of climate and its
1310 interaction with species-level functional trait data (WD and SLA). This is based on a subset
1311 of the mortality data, for which trait information was available (a). Parameter estimates for
1312 each of the fixed effects included in the best trait model, with 95% confidence intervals,
1313 calculated using bootstrapped resampling (b).

1314

1315 Figure 6: The effect of the interaction of mean SPEI of the reported drought years with wood
1316 density (WD) (a) and specific leaf area (SLA) (b) on log drought-induced tree mortality (trees
1317 $\text{trees}^{-1} \text{ year}^{-1}$).

1318




Figure 1






Figure 2



1323

1324 **Figure 3**



1325

1326 **Figure 4**

1327




Figure 5




Figure 6